

# SHARP

## Werkkaart 11 Memorandum: Calculus Deel 2

### Graad 12 Wiskunde

1. a)  $y = x^2 + 3x - 12$   
 $\frac{dy}{dx} = 2x + 3$   
 $\frac{d^2y}{dx^2} = 2$
- b)  $f(x) = 4x - 5$   
 $f'(x) = 4$   
 $f''(x) = 0$
- c)  $y = x^3 - 4x^2 + 8x - 7$   
 $\frac{dy}{dx} = 3x^2 - 8x + 8$   
 $\frac{d^2y}{dx^2} = 6x - 8$
- d)  $y = \sqrt[3]{x} + \frac{1}{x^2}$   
 $y = x^{\frac{1}{3}} + x^{-2}$   
 $\frac{dy}{dx} = \frac{1}{3}x^{-\frac{2}{3}} - 2x^{-3} = \frac{1}{3\sqrt[3]{x^2}} - \frac{2}{x^3}$   
 $\frac{d^2y}{dx^2} = -\frac{2}{9}x^{-\frac{5}{3}} + 6x^{-4} = -\frac{2}{9\sqrt[3]{x^5}} + \frac{6}{x^4}$
- e)  $f(x) = \frac{5}{x} - 11x^2$   
 $f(x) = 5x^{-1} - 11x^2$   
 $f'(x) = -5x^{-2} - 22x = -\frac{5}{x^2} - 22x$   
 $f''(x) = 10x^{-3} - 22 = \frac{10}{x^3} - 22$
- f)  $g(x) = \frac{x^2 - x}{x - 1}$   
 $g(x) = \frac{x(x-1)}{(x-1)} = x$   
 $g'(x) = 1$   
 $g''(x) = 0$
- g)  $y = x^5 + 7x^2 - 12$   
 $\frac{dy}{dx} = 5x^4 + 14x$   
 $\frac{d^2y}{dx^2} = 20x^3 + 14$
- h)  $y = -x^3 + 4x^2 - 18x - 2$   
 $\frac{dy}{dx} = -3x^2 + 8x - 18$   
 $\frac{d^2y}{dx^2} = -6x + 8$
- i)  $h(x) = \frac{4}{x^3} - \sqrt{x^3}$   
 $h(x) = 4x^{-3} - x^{\frac{3}{2}}$   
 $h'(x) = -12x^{-4} - \frac{3}{2}x^{\frac{1}{2}} = -\frac{12}{x^4} - \frac{3}{2}\sqrt{x}$   
 $h''(x) = 48x^{-5} - \frac{3}{4}x^{-\frac{1}{2}} = \frac{48}{x^5} - \frac{3}{4\sqrt{x}}$
- j)  $y = \frac{1}{3}x^3 + \frac{1}{4}x^2 - 3x + 2$   
 $\frac{dy}{dx} = x^2 + \frac{1}{2}x - 3$   
 $\frac{d^2y}{dx^2} = 2x + \frac{1}{2}$
2. a)  $f(x) = x^3 - 5x^2 - 10x + 5$   
 $f'(x) = 3x^2 - 10x - 10$   
 $f''(x) = 6x - 10$   
Om die buigpunt te vind, stel die tweede afgeleide gelyk aan nul:  
 $f''(x) = 6x - 10 = 0$   
 $6x = 10$   
 $x = \frac{10}{6}$  of  $\frac{5}{3}$
- b)  $g(x) = -x^3 + x^2 - 3x + 7$   
 $g'(x) = -3x^2 + 2x - 3$   
 $g''(x) = -6x + 2$   
 $g''(x) = -6x + 2 = 0$   
 $-6x = -2$   
 $x = \frac{1}{3}$

$$\text{En } y = \left(\frac{5}{3}\right)^3 - 5\left(\frac{5}{3}\right)^2 - 10\left(\frac{5}{3}\right) + 5$$

$$y = -20\frac{25}{27}$$

$$\therefore \text{ buigpunt: } \left(\frac{5}{3}; -20\frac{25}{27}\right)$$

Omdat die grafiek 'n algemeen toenemende grafiek is, is die grafiek

Konkaaf af vir  $x < \frac{5}{3}$ , en

konkaaf op vir  $x > \frac{5}{3}$

$$\text{En } y = -\left(\frac{1}{3}\right)^3 + \left(\frac{1}{3}\right)^2 - 3\left(\frac{1}{3}\right) + 7$$

$$y = 6\frac{2}{27}$$

$$\therefore \text{ buigpunt: } \left(\frac{1}{3}; 6\frac{2}{27}\right)$$

Omdat die grafiek 'n algemeen

afnemende grafiek, is die grafiek konkaaf

op vir  $x < \frac{1}{3}$ , en konkaaf af vir  $x > \frac{1}{3}$ .

c)  $h(x) = (x - 3)(x + 2)(x - 1)$

$$h(x) = x^3 - 2x^2 - 5x + 6$$

$$h'(x) = 3x^2 - 4x - 5$$

$$h''(x) = 6x - 4 = 0$$

$$\therefore x = \frac{4}{6} \text{ or } \frac{2}{3}$$

$$y = \left(\frac{2}{3}\right)^3 - 2\left(\frac{2}{3}\right)^2 - 5\left(\frac{2}{3}\right) + 6$$

$$y = 2\frac{2}{27}$$

$$\therefore \text{ buigpunt is } \left(\frac{2}{3}; 2\frac{2}{27}\right)$$

Omdat die grafiek algemeen

toenemende grafiek is, is die grafiek

konkaaf af vir  $x < \frac{2}{3}$ , en

konkaaf op vir  $x > \frac{2}{3}$

d)  $j(x) = x^3 - 2$

$$j'(x) = 3x^2$$

$$j''(x) = 6x = 0$$

$$\therefore x = 0$$

$$y = 0^3 - 2$$

$$y = -2$$

$$\therefore \text{ buigpunt is } (0; -2)$$

Omdat die grafiek 'n algemeen

toenemende grafiek is, die grafiek is

konkaaf af vir  $x < 0$ , en konkaaf op vir  $x > 0$

e)  $k(x) = -3x^3 + 12x^2$

$$k'(x) = -9x^2 + 24x$$

$$k''(x) = -18x + 24 = 0$$

$$x = \frac{4}{3}$$

$$\therefore y = -3\left(\frac{4}{3}\right)^3 + 12\left(\frac{4}{3}\right)^2$$

$$\therefore y = 14\frac{2}{9}$$

$$\therefore \text{ die buigpunt is } \left(\frac{4}{3}; 14\frac{2}{9}\right)$$

Omdat die grafiek 'n algemeen afnemende grafiek is, is die grafiek konkaaf op vir  $x <$

$\frac{4}{3}$ , en konkaaf af vir  $x > \frac{4}{3}$ .

f)  $m(x) = x^2 - 3x + 12$

$$m'(x) = 2x - 3$$

$$m''(x) = 2$$

Die grafiek het nie 'n buiging nie

punt en is konkaaf af oor die geheel grafiek.

g)  $n(x) = -(x - 1)(x - 2)(x - 4)$

$$n(x) = -x^3 + 7x^2 - 14x + 8$$

$$n'(x) = -3x^2 + 14x - 14$$

$$n''(x) = -6x + 14 = 0$$

$$x = \frac{7}{3}$$

$$y = -\left(\frac{7}{3}\right)^3 + 7\left(\frac{7}{3}\right)^2 - 14\left(\frac{7}{3}\right) + 8$$

$$y = \frac{20}{27}$$

$$\therefore \text{ buigpunt by } \left(\frac{7}{3}; \frac{20}{27}\right)$$

h)  $p(x) = -(x^2 - 8)(x + 3)$

$$p(x) = -x^3 - 3x^2 + 8x + 24$$

$$p'(x) = -3x^2 - 6x + 8$$

$$p''(x) = -6x - 6 = 0$$

$$x = -1$$

$$y = -(-1)^3 - 3(-1)^2 + 8(-1) + 24$$

$$y = 14$$

$$\therefore \text{ buigpunt by } (-1; 14)$$

Omdat die grafiek 'n algemeen afnemende grafiek is, is die grafiek konkaf op vir  $x < \frac{7}{3}$ , en konkaf af vir  $x > \frac{7}{3}$

Omdat die grafiek 'n algemeen afnemende grafiek, die grafiek is konkaf op vir  $x < -1$ , en konkaf af vir  $x > -1$

i)  $q(x) = 4x^3 - 6x^2 - 3x + 5$   
 $q'(x) = 12x^2 - 12x - 3$   
 $q''(x) = 24x - 12 = 0$   
 $x = \frac{1}{2}$

$$y = 4\left(\frac{1}{2}\right)^3 - 6\left(\frac{1}{2}\right)^2 - 3\left(\frac{1}{2}\right) + 5$$

$$y = 2\frac{1}{2}$$

$\therefore$  buigpunt  $\left(\frac{1}{2}; 2\frac{1}{2}\right)$

Omdat die grafiek 'n algemeen toenemende grafiek is, is die grafiek konkaf af vir  $x < \frac{1}{2}$ , en konkaf op vir  $x > \frac{1}{2}$

j)  $r(x) = 5x^3 + 3x^2 - 4x + 12$   
 $r'(x) = 15x^2 + 6x - 4$   
 $r''(x) = 30x + 6 = 0$   
 $x = -\frac{1}{5}$

$$y = 5\left(-\frac{1}{5}\right)^3 + 3\left(-\frac{1}{5}\right)^2 - 4\left(-\frac{1}{5}\right) + 12$$

$$y = 12\frac{22}{25} \text{ or } \frac{322}{25}$$

$\therefore$  buigpunt  $\left(-\frac{1}{5}; 12\frac{22}{25}\right)$

Omdat die grafiek 'n algemeen konkaf toenemende grafiek, die grafiek af vir  $x < -\frac{1}{5}$ , en konkaf op vir  $x > -\frac{1}{5}$

3. a)  $y = x^3 + 7x^2 + 7x - 15$

i) y-afsnit  $(0; -15)$   
x-afsnit:

$$0 = x^3 + 7x^2 + 7x - 15 \quad x = 1 \text{ of } (x - 1) \text{ is 'n faktor:}$$

$$0 = (x - 1)(x^2 + kx + 15)$$

Om k te vind:

$$-x^2 + kx^2 = 7x^2$$

$$kx^2 = 8x^2$$

$$\therefore k = 8$$

$$0 = (x - 1)(x^2 + 8x + 15)$$

$$0 = (x - 1)(x + 3)(x + 5)$$

$\therefore$  die x-afsnitte is  $(1; 0)$ ,  $(-3; 0)$  en  $(-5; 0)$

ii)  $f'(x) = 3x^2 + 14x + 7 = 0$

$$x = \frac{-14 \pm \sqrt{14^2 - 4(3)(7)}}{2(3)}$$

$$x = \frac{-7 \pm 2\sqrt{7}}{3}$$

$$x = -0.57 \quad \text{of} \quad x = -4.1$$

$$y = (-0.57)^3 + 7(-0.57)^2 + 7(-0.57) - 15 = -16.9$$

$$\text{En } y = (-4.1)^3 + 7(-4.1)^2 + 7(-4.1) - 15 = 5.05$$


dus :  $(-4.1; 5.05)$  en  $(-0.57; -16.9)$

iii)  $f''(x) = 6x + 14 = 0$

$$\therefore x = -\frac{14}{6} \text{ of } -2\frac{1}{3}$$

$$\therefore y = \left(-2\frac{1}{3}\right)^3 + 7\left(-2\frac{1}{3}\right)^2 + 7\left(-2\frac{1}{3}\right) - 15$$

$$y = -5\frac{25}{27}$$



iv)

b)  $y = x^3 + 13x^2 + 34x - 48$

i) y-afsnit: (0; -48)

Vir die x-afsnitte:

$$0 = x^3 + 13x^2 + 34x - 48$$

(x - 1) is 'n faktor

$$\therefore (x - 1)(x^2 + kx + 48)$$

$$-x^2 + kx^2 = 13x^2$$

$$kx^2 = 14x^2$$

$$\therefore k = 14$$

$$\therefore 0 = (x - 1)(x^2 + 14x + 48)$$

$$0 = (x - 1)(x + 6)(x + 8)$$

$\therefore$  Die x-afsnitte is: (1; 0), (-6; 0) en (-8; 0)

ii)  $f'(x) = 3x^2 + 26x + 34 = 0$

$$x = \frac{-26 \pm \sqrt{26^2 - 4(3)(34)}}{2(3)}$$

$$x = \frac{-13 \pm \sqrt{67}}{3}$$

$$x = -1.6 \text{ of } x = -7.1$$

$$\therefore y = (-1.6)^3 + 13(-1.6)^2 + 34(-1.6) - 48 = -73.2$$


$$\text{En } y = (-7.1)^3 + 13(-7.1)^2 + 34(-7.1) - 48 = 8$$

iii)  $f''(x) = 6x + 26 = 0$

$$\therefore x = -4\frac{1}{3}$$

$$\therefore y = \left(-4\frac{1}{3}\right)^3 + 13\left(-4\frac{1}{3}\right)^2 + 34\left(-4\frac{1}{3}\right) - 48 = -32\frac{16}{27}$$

$\therefore$  die buigpunt is  $\left(-4\frac{1}{3}; -32\frac{16}{27}\right)$



iv)

c)  $y = 3x^3 - 13x^2 - 130x + 336$

i) y-afsnit:  $(0; 336)$

x-afsnitte:

$$0 = 3x^3 - 13x^2 - 130x + 336 \quad (x - 8) \text{ is 'n faktor}$$

$$\therefore 0 = (x - 8)(3x^2 + kx - 42)$$

$$kx^2 - 24x^2 = -13x^2$$

$$kx^2 = 11x^2$$

$$\therefore k = 11$$

$$\therefore 0 = (x - 8)(3x^2 + 11x - 42)$$

$$\therefore 0 = (x - 8)(3x - 7)(x + 6)$$

$$\therefore \text{die x-afsnitte is: } (8; 0), (-6; 0) \text{ en } \left(2\frac{1}{3}; 0\right)$$

ii)  $\frac{dy}{dx} = 9x^2 - 26x - 130 = 0$

$$x = \frac{26 \pm \sqrt{(-26)^2 - 4(9)(-130)}}{2(9)}$$

$$x = 5.51 \quad \text{of} \quad x = -2.62$$

$$\therefore y = 3(5.51)^3 - 13(5.51)^2 - 130(5.51) + 336 = -273.13$$

$$\text{En } y = 3(-2.62)^3 - 13(-2.62)^2 - 130(-2.62) + 336 = 533.4$$


$$\therefore \text{die maksimum en minimum is: } (-2.62; 533.4) \text{ en } (5.51; -273.13)$$

iii)  $\frac{d^2y}{dx^2} = 18x - 26 = 0$

$$x = 1\frac{4}{9}$$

$$\therefore y = 3\left(1\frac{4}{9}\right)^3 - 13\left(1\frac{4}{9}\right)^2 - 130\left(1\frac{4}{9}\right) + 336 = 130.14$$

$$\therefore \text{buigpunt: } \left(1\frac{4}{9}; 130.14\right)$$



iv)

d)  $y = -x^3 + 6x^2 + 45x - 162$

i) y-afsnit:  $(0; -162)$

Vir die x-afsnitte:

$$0 = -(x^3 - 6x^2 - 45x + 162) \quad \therefore (x - 3) \text{ is faktor}$$

$$0 = -(x - 3)(x^2 + kx - 54)$$

$$kx^2 - 3x^2 = -6x^2$$

$$kx^2 = -3x^2$$

$$k = -3$$

$$\therefore 0 = -(x - 3)(x^2 - 3x - 54)$$

$$\therefore 0 = -(x - 3)(x + 6)(x - 9)$$

$\therefore$  Die x-afsnitte is:  $(3; 0)$ ,  $(-6; 0)$  en  $(9; 0)$

ii)  $\frac{dy}{dx} = -3x^2 + 12x + 45 = 0$

$$x^2 - 4x - 15 = 0$$

$$x = \frac{4 \pm \sqrt{(-4)^2 - 4(1)(-15)}}{2}$$

$$x = 2 \pm \sqrt{19}$$

$$x = 6.36 \quad \text{or} \quad x = -2.36$$

$$\therefore y = -(6.36)^3 + 6(6.36)^2 + 45(6.36) - 162 = 109.64$$

$$\text{En } y = -(-2.36)^3 + 6(-2.36)^2 + 45(-2.36) - 162 = -221.64$$


$\therefore$  die maksimum en minimum is:  $(6.36; 109.64)$  en  $(-2.36; -221.64)$

iii)  $\frac{d^2y}{dx^2} = -6x + 12 = 0$

$$x = 2$$

$$\therefore y = -(2)^3 + 6(2)^2 + 45(2) - 162 = -56$$

$\therefore$  die buigpunt is  $(2; -56)$ .



iv)

e)  $y = 3x^3 - 29x^2 + 17x + 9$

i) die y-afsnit is (0; 9)

Vir die x-afsnitte:

$$0 = 3x^3 - 29x^2 + 17x + 9 \quad (x - 1) \text{ is 'n faktor}$$

$$\therefore 0 = (x - 1)(3x^2 + kx - 9)$$

$$kx^2 - 3x^2 = -29x^2$$

$$kx^2 = -26x^2$$

$$\therefore k = -26$$

$$\therefore 0 = (x - 1)(3x^2 - 26x - 9)$$

$$\therefore 0 = (x - 1)(3x + 1)(x - 9)$$

$\therefore$  Die x-afsnitte is: (1; 0), (9; 0) en  $(-\frac{1}{3}; 0)$

ii)  $\frac{dy}{dx} = 9x^2 - 58x + 17 = 0$

$$x = \frac{58 \pm \sqrt{(-58)^2 - 4(9)(17)}}{2(9)}$$

$$x = \frac{29 \pm 4\sqrt{43}}{9}$$

$$x = 6.14 \quad \text{of} \quad x = 0.31$$

$$\therefore y = 3(6.14)^3 - 29(6.14)^2 + 17(6.14) + 9 = -285.48$$

$$\text{En } y = 3(0.31)^3 - 29(0.31)^2 + 17(0.31) + 9 = 11.57$$


$\therefore$  die maksimum en minimum is: (0.31; 11.57) en (6.14; -285.48)

iii)  $\frac{d^2y}{dx^2} = 18x - 58 = 0$

$$x = 3\frac{2}{9}$$

$$\therefore y = 3\left(3\frac{2}{9}\right)^3 - 29\left(3\frac{2}{9}\right)^2 + 17\left(3\frac{2}{9}\right) + 9 = -136.95$$

$\therefore$  die buigpunt is  $\left(3\frac{2}{9}; -136.95\right)$



iv)

f)  $y = -3x^3 + 20x^2 + 69x - 54$

i) y-afsnit:  $(0; -54)$

Vir die x-afsnitte:

$$0 = -(3x^3 - 20x^2 - 69x + 54) \quad (x - 9) \text{ is 'n faktor}$$

$$\therefore 0 = -(x - 9)(3x^2 + kx - 6)$$

$$kx^2 - 27x^2 = -20x^2$$

$$kx^2 = 7x^2$$

$$\therefore k = 7$$

$$\therefore 0 = -(x - 9)(3x^2 + 7x - 6)$$

$$\therefore 0 = -(x - 9)(3x - 2)(x + 3)$$

$$\therefore \text{die x-afsnitte is: } (9; 0), (-3; 0) \text{ en } \left(\frac{2}{3}; 0\right)$$

ii)  $\frac{dy}{dx} = -9x^2 + 40x + 69 = 0$

$$x = \frac{-40 \pm \sqrt{40^2 - 4(-9)(69)}}{2(-9)}$$

$$x = -1.33 \quad \text{or} \quad x = 5.77$$

$$\therefore y = -3(-1.33)^3 + 20(-1.33)^2 + 69(-1.33) - 54 = -103.33$$

$$\text{En } y = -3(5.77)^3 + 20(5.77)^2 + 69(5.77) - 54 = 433.69$$

$$\therefore \text{die maksimum en die minimum is } (5.77; 433.69) \text{ en } (-1.33; -103.33)$$


iii)  $\frac{d^2y}{dx^2} = -18x + 40 = 0$

$$x = 2\frac{2}{9}$$

$$\therefore y = -3\left(2\frac{2}{9}\right)^3 + 20\left(2\frac{2}{9}\right)^2 + 69\left(2\frac{2}{9}\right) - 54 = 165.18$$

$$\therefore \text{die buigpunt is } \left(2\frac{2}{9}; 165.18\right)$$





iv)

g)  $y = 5x^3 - 39x^2 + 78x - 40$

i) y-afsnit:  $(0; -40)$

Vir die x-afsnitte:

$$0 = 5x^3 - 39x^2 + 78x - 40$$

$(x - 2)$  is 'n faktor

$$\therefore 0 = (x - 2)(5x^2 + kx + 20)$$

$$kx^2 - 10x^2 = -39x^2$$

$$kx^2 = -29x^2$$

$$\therefore k = -29$$

$$\therefore 0 = (x - 2)(5x^2 - 29 + 20)$$

$$\therefore 0 = (x - 2)(5x - 4)(x - 5)$$

$$\therefore \text{die x-afsnitte is: } (2; 0), (5; 0) \text{ en } \left(\frac{4}{5}; 0\right)$$

ii)  $\frac{dy}{dx} = 15x^2 - 78x + 78 = 0$

$$x = \frac{78 \pm \sqrt{(-78)^2 - 4(15)(78)}}{2(15)}$$

$$x = \frac{13 \pm \sqrt{39}}{5}$$


$$x = 3.85 \quad \text{of} \quad x = 1.35$$

$$\therefore y = 5(3.85)^3 - 39(3.85)^2 + 78(3.85) - 40 = -32.44$$

$$\text{En } y = 5(1.35)^3 - 39(1.35)^2 + 78(1.35) - 40 = 6.52$$

$$\therefore \text{die maksimum en minimum is } (1.35; 6.52) \text{ en } (3.85; -32.44)$$

iii)  $\frac{d^2y}{dx^2} = 30x - 78 = 0$   
 $x = 2\frac{3}{5}$   
 $\therefore y = 5\left(2\frac{3}{5}\right)^3 - 39\left(2\frac{3}{5}\right)^2 + 78\left(2\frac{3}{5}\right) - 40 = -12.96$   
 $\therefore$  die buigpunt is  $\left(2\frac{3}{5}; -12.96\right)$



h)  $y = -x^3 + 6x^2 + 67x - 360$

i) y-afsnit:  $(0; -360)$

Vir die x-afsnitte:

$0 = -(x^3 - 6x^2 - 67x + 360)$        $(x - 5)$  is 'n faktor

$\therefore 0 = -(x - 5)(x^2 + kx - 72)$

$kx^2 - 5x^2 = -6x^2$

$kx^2 = -x^2$

$\therefore k = -1$

$\therefore 0 = -(x - 5)(x - x - 72)$

$\therefore 0 = -(x - 5)(x - 9)(x + 8)$

$\therefore$  die x-afsnitte is  $(5; 0)$ ,  $(9; 0)$  en  $(-8; 0)$

ii)  $\frac{dy}{dx} = -3x^2 + 12x + 67 = 0$

$x = \frac{-12 \pm \sqrt{12^2 - 4(-3)(67)}}{2(-3)}$

$x = \frac{6 \mp \sqrt{237}}{3}$

$$x = -3.13 \quad \text{of} \quad x = 7.13$$

$$\therefore y = -(-3.13)^3 + 6(-3.13)^2 + 67(-3.13) - 360 = -480.26$$

$$\text{En } y = -(7.13)^3 + 6(7.13)^2 + 67(7.13) - 360 = 60.26$$


$$\therefore \text{die maksimum en minimum is: } (7.13; 60.26) \text{ en } (-3.13; -480.26)$$

iii)  $\frac{d^2y}{dx^2} = -6x + 12 = 0$

$$x = 2$$

$$\therefore y = -(2)^3 + 6(2)^2 + 67(2) - 360 = -210$$

$$\therefore \text{die buigpunt is } (2; -210)$$



ek)  $y = 6x^3 + 7x^2 - 84x - 160$

i) y-afsnit:  $(0; -160)$   
Vir die x-afsnitte:

$$0 = 6x^3 + 7x^2 - 84x - 160 \quad (x - 4) \text{ is 'n faktor}$$

$$\therefore 0 = (x - 4)(6x^2 + kx + 40)$$

$$kx^2 - 24x^2 = 7x^2$$

$$kx^2 = 31x^2$$

$$\therefore k = 31$$

$$\therefore 0 = (x - 4)(6x^2 + 31x + 40)$$

$$\therefore 0 = (x - 4)(3x + 8)(2x + 5)$$

$$\therefore \text{die x-afsnitte is: } (4; 0), \left(-2\frac{2}{3}; 0\right) \text{ en } \left(-2\frac{1}{2}; 0\right)$$

ii)  $\frac{dy}{dx} = 18x^2 + 14x - 84 = 0$

$$x = \frac{-14 \pm \sqrt{14^2 - 4(18)(-84)}}{2(18)}$$

$$x = 1.81 \quad \text{of} \quad x = -2.58$$


$$\begin{aligned} \therefore y &= 6(1.81)^3 + 7(1.81)^2 - 84(1.81) - 160 = -253.5 \\ \text{En } y &= 6(-2.58)^3 + 7(-2.58)^2 - 84(-2.58) - 160 = 0.27 \\ \therefore \text{die maksimum en minimum is: } &(-2.58; 0.27) \text{ en } (1.81; -253.5) \end{aligned}$$

iii)  $\frac{d^2y}{dx^2} = 36x + 14 = 0$

$$x = \frac{7}{18}$$

$$\therefore y = 6\left(\frac{7}{18}\right)^3 + 7\left(\frac{7}{18}\right)^2 - 84\left(\frac{7}{18}\right) - 160 = -191.26$$

$\therefore$  die buigpunt is  $\left(\frac{7}{18}; -191.26\right)$



iv)

j)  $y = -x^3 - 7x^2 + 84x + 288$

i) y-afsnit:  $(0; 288)$

Vir die x-afsnitte:

$$0 = -(x^3 + 7x^2 - 84x - 288) \quad (x - 8) \text{ is 'n faktor}$$

$$\therefore 0 = -(x - 8)(x^2 + kx + 36)$$

$$kx^2 - 8x^2 = 7x^2$$

$$kx^2 = 15x^2$$

$$\therefore k = 15$$

$$\therefore 0 = -(x - 8)(x^2 + 15x + 36)$$

$$\therefore 0 = -(x - 8)(x + 12)(x + 3)$$

$$\therefore \text{die x-afsnitte is: } (8; 0), (-12; 0) \text{ en } (-3; 0)$$

ii)  $\frac{dy}{dx} = -3x^2 - 14x + 84 = 0$

$$x = \frac{14 \pm \sqrt{(-14)^2 - 4(-3)(84)}}{2(-3)}$$

$$x = \frac{-7 \mp \sqrt{301}}{3}$$

$x = -8.12$  of  $x = 3.45$

$\therefore y = -(-8.12)^3 - 7(-8.12)^2 + 84(-8.12) + 288 = -320.23$


En  $y = -(3.45)^3 - 7(3.45)^2 + 84(3.45) + 288 = 453.42$

$\therefore$  die maksimum en minimum is: (3.45; 453.42) en (-8.12; -320.23)

iii)  $\frac{d^2y}{dx^2} = -6x - 14 = 0$

$$x = -2\frac{1}{3}$$

$\therefore y = -\left(-2\frac{1}{3}\right)^3 - 7\left(-2\frac{1}{3}\right)^2 + 84\left(-2\frac{1}{3}\right) + 288 = 66.59$



iv)

4. a)  $v(t) = \frac{1}{3}t^3 - 3t^2 + 6t = 0$

$$\frac{1}{3}t(t^2 - 9t + 18) = 0$$

$$\frac{1}{3}t(t-3)(t-6) = 0$$

$\therefore t = 0, t = 3$  or  $t = 6$

Die molekule kan nie deur die selwand kom as die grafiek negatief is nie, en dit is tussen  $t = 3$  en  $t = 6$

b)  $v'(t) = t^2 - 6t + 6 = 0$

$$t = \frac{6 \pm \sqrt{(-6)^2 - 4(1)(6)}}{2}$$

$$t = 3 \pm \sqrt{3}$$

$t = 4.73$  of  $t = 1.27$

$$\therefore v(4.73) = \frac{1}{3}(4.73)^3 - 3(4.73)^2 + 6(4.73) = -3.46$$

$$\text{En } v(1.27) = \frac{1}{3}(1.27)^3 - 3(1.27)^2 + 6(1.27) = 3.46$$

Daarom is die maksimum punt by (1.27; 3.46)

5. Volume van oorspronklike boks =  $x^2 y = 3$

$$\therefore y = \frac{3}{x^2}$$

Buite-Oppervlakte van die boks =  $2x^2 + 2xy + 2xy$

$$BO = 2x^2 + 4xy \quad \text{Vervang volume vir } y:$$

$$BO = 2x^2 + 4x \left( \frac{3}{x^2} \right)$$

$$BO = 2x^2 + \frac{12}{x} = 2x^2 + 12x^{-1}$$

Differensieer die buit-oppervlakte en stel gelyk aan nul:

$$SA' = 4x - 12x^{-2} = 0$$

$$4x - \frac{12}{x^2} = 0$$

$$4x^3 - 12 = 0$$

$$4x^3 = 12$$

$$x^3 = 3$$

$$x = \sqrt[3]{3} \approx 1.44$$

$$\therefore \text{die maksimum oppervlakte is } BO = 2(1.44)^2 + \frac{12}{1.44} = 12.48 \text{ units}^2$$

6. a)  $P = 300m = 2x + 2y$

$$2y = 300 - 2x$$

$$y = 150 - x$$

- b)  $A = xy$

$$A = x(150 - x)$$

$$A = 150x - x^2$$

$$\therefore A' = 150 - 2x = 0$$

$$x = 75m$$

$$\therefore \text{die maksimum oppervlakte is } A = 150(75) - (75)^2 = 5\,625m^2$$

- c)  $\therefore$  nuwe omtrek = 380m

$$2y = 380 - 2x$$

$$y = 190 - x$$

- d)  $A = x(190 - x)$

$$A = 190x - x^2$$

$$\therefore A' = 190 - 2x = 0$$

$$x = 95m$$

$$\therefore \text{die nuwe maksimum area is } A = 190(95) - (95)^2 = 9\,025m^2$$